

UNCLASSIFIED

AD 4 3 9 2 9 5

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

DDC
CATALOG
AS AD NO.

439295

439295

Quarterly REPORT ON

CONTRACT NO DA - 92 - 557 - FEC -35580

INCLUSIVE DATES 1 April 1963 TO 30 June 1963

SUBJECT OF INVESTIGATION

BACTERIOLOGICAL, IMMUNOLOGICAL
AND
VIRAL STUDIES ON RECTAL MUCUS
IN
ENTERIC INFECTIONS

(SHIGELLOSIS, SALMONELLOSIS,
PATHOGENIC COLI INFECTIONS
AND VIRAL ENTERIC INFECTIONS)

RESPONSIBLE INVESTIGATOR

Dr. Yasuzo Nakamizo

Chief Doctor
Tokyo Metropolitan Ebara
Infectious Disease Hospital
895, Yurigaya, Ota-ku, Tokyo, Japan

MAY 12 1964

U.S. Army Research & Development Group (9852) (Far East)

Office of the Chief of Research and Development

United States Army

APO 343

64-13

J-177

D-I-S-T-R-I-B-U-T-I-O-N

The distribution of this report as made by USA R&D Gp (FE)
is as follows:

Army Research Office, 3045 Columbia Pike, Arlington 4, Virginia. ATTN: Chief Research Programs Office (h) /

Army Attache, American Embassy, Tokyo, Japan (1)

Office of Primary Scientific Liaison (h)
U.S. Army Medical Research & Development Command
ATTN: Chief, Preventive Medicine Research Br
Washington 25, D. C.

Offices of Scientific Cognizance (1)
Chief Applied Immunology,
Division of Communicable Diseases, WRAIR
ATTN: Dr. S. B. Fernal
Washington 12, D. C.

BACTERIOLOGICAL, IMMUNOLOGICAL AND
VIRAL STUDIES ON RECTAL MUCUS IN
ENTERIC INFECTIONS
(SHIGELLOSIS, SALMONELLOSIS, PATHO-
GENIC COLI INFECTIONS AND VIRAL
ENTERIC INFECTIONS)

Serological findings on serum and rectal mucus in acute gastro-enteritis patients and Salmonellosis, viral researches on feces and mucus in suspected cases of viral enteric infections obtained during the period of 1 April 1963 through 30 June 1963 were summarized as follows:

1. Agglutinin titers of serum and mucus against O-antigen and H-antigen of six entero-bacilli (Morganella 1, A. liquefaciens 1, Escherichia coli 4) isolated from feces of acute gastro-enteritis patients were very low as seen in Table 1.

Although clinical figures of these cases were similar to bacillary dysentery, no Shigella strain isolated.

2. Comparative study of agglutinin titers between the Widal test and the Latex Widal test in sera of twenty-one cases of typhoid fever patients was carried out. The Latex agglutinin titers were mostly higher than the Widal agglutinin titers in both agglutinin titers against Vi-antigen or O-antigen of Salmonella typhi. (Table 2)

3. Influences of three solutions (0.5% saline, 0.85% saline, glycine saline buffer Difco) and Latex solution upon agglutinin titer of rabbit anti-Salmonella typhi serum and antigenicity of O-antigen of Salmonella typhi were studied.

It was clarified that glycine saline buffer Difco is most suitable to detection of O-agglutinin of anti-O serum. (Table 3)

4. Horie reported at the meeting of Society of Japanese Virologists 1962 that isolation and differentiation of entero-viruses were demonstrated using his four tissue culture cells.

He pointed out regular occurrences of cytopathic effects of entero-viruses in four cell lines as shown in Table 4.

Monolayer cultures of these tissue culture cells were grown in growth medium consisted of 0.1% yeast extract, 0.5% lactalbumin hydrolysate in Earle solution with 20% calf serum.

Our trial for detecting entero-virus from feces and rectal mucus of suspected ten cases using Horie's four cells fell through.

Table 1. Agglutinin titers of serum and mucus against O-antigen and H-antigen of entero-bacilli isolated from feces of acute gastro-enteritis patients

No.	Entero-bacilli isolated	Week of illness	Material	Agglutinin titer against O-antigen	Agglutinin titer against H-antigen
1	Morganella	3	serum mucus	< 20 < 20	< 20 < 20
2	A. liquefaciens	3	serum mucus	< 20 < 20	< 20 < 20
3	Escherichia coli	3	serum mucus	< 20 < 20	< 20 < 20
4	Escherichia coli	3	serum mucus	< 20 < 20	40 < 20
5	Escherichia coli	2	serum mucus	< 20 < 20	< 20 < 20
6	Escherichia coli	2	serum mucus	< 20 < 20	< 20 < 20

Table 2. Comparative study of the Widal test and the Latex Widal test in sera of typhoid fever patients

Widal test antigen	Agglutinin titer	Widal test	Latex Widal test
Vi-antigen	< 10 x	6	5
	10 x	6	6
	20 x	3	2
	40 x	6	7
	80 x	0	1
O-antigen of Salmonella typhi	< 40 x	1	2
	40 x	4	1
	80 x	8	4
	160 x	6	9
	320 x	2	5

Table 3. Influences of three saline solutions and Latex solution upon agglutinin titer of rabbit anti-O 901 serum and antigenicity of Salmonella typhi

Saline solution for diluent of rabbit anti-O 901 serum	Saline solution for diluent of O 901 strain concentration of 1 mg per 1 ml	Latex suspension (10x) 0.1 ml to 10 ml antigen solution	Agglutinin titer
0.5%	0.5%	-	5120 x
0.5%	0.5%	+	5120 x
0.5%	0.85%	-	5120 x
0.5%	0.85%	+	5120 x
0.5%	G.S.B.	-	5120 x
0.5%	G.S.B.	+	10240 x
0.85%	0.5%	-	2560 x
0.85%	0.5%	+	5120 x
0.85%	0.85%	-	5120 x
0.85%	0.85%	+	5120 x
0.85%	G.S.B.	-	10240 x
0.85%	G.S.B.	+	10240 x
G.S.B.	0.5%	-	10240 x
G.S.B.	0.5%	+	10240 x
G.S.B.	0.85%	-	10240 x
G.S.B.	0.85%	+	10240 x
G.S.B.	G.S.B.	-	10240 x
G.S.B.	G.S.B.	+	10240 x

* Agglutination was observed after incubation for 90 M. at 50 C and centrifugation for 5 M. at 2500 r.p.m.

* G.S.B. Glycine saline buffer Difco

Table 4. Regular occurrences of cytopathic effects of enteroviruses in Horie's four tissue culture cells

Tissue culture cell	Polio-virus	Coxsackie-virus	ECHO-virus
FL-17-M	CPE	CPE	CPE
1035-1	-	CPE	CPE
E 10	-	CPE	-
E 38	CPE	CPE	-

* CPE cytopathic effect